

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

accurate position can be found if photographic charts can be taken showing this star. Professor Bailey has been notified of this discovery and if the star is still bright enough, he will doubtless obtain photographs showing its position and spectrum.

"The similarity of these two new stars is interesting; first, since it has proved a means of discovering one of these objects, and secondly, because if confirmed by other new stars it will indicate that they belong to a distinct class resembling each other in composition or physical condition.

"The star was approximately of the 7th magnitude photographically on July 10, since it was about equal to *Cord. G. C.* 20910, magnitude 6.9, whose spectrum is of the second type. The nearest catalogue stars are *Cord. G. C.* 20940, magnitude 8, which has a spectrum of the first type, and *Cord. G. C.* 20926, magnitude $8\frac{3}{4}$, which has a spectrum of the second type. The new star lies nearly midway between these two."

EDWARD C. PICKERING,
Director Harvard College Observatory.
CAMBRIDGE, MASS., U. S. A., Nov. 9, 1893.

THE DISCOVERY OF ASTEROIDS IN 1893.

Thirty-four minor planets were discovered in 1893: eight by Wolf of Heidelberg, twenty-five by Charlos of Nice and one by Borelly of Marseilles. Wolf and Charlos used the photographic method, first successfully employed by Wolf in 1891.

The number of minor planets now known is about 380. The rate of their discovery by photography in the past two years has been so rapid that the computers are no longer able to make the computations for determining their orbits properly. We believe that Dr. Wolf has for this reason desisted from making further special search for them. Dr. Wolf's decision is to be commended. The cosmical questions with which the asteroids may be connected will certainly be nearest solution if the computations for those already discovered are kept up to date.

W. W. C.

Conferring of the Degree of Doctor of Mathematical Sciences upon Miss Dorothea Klumpke.

Mlle. Klumpke, who has just gained the degree of Doctor of Mathematical Sciences at the Sorbonne, is the first lady who has obtained that distinction. The full title of her thesis was "Con-